Application of Radar Remote Sensing in Mapping and Monitoring Tropical Forest Types in Amazon Basin

Sassan Saatchi¹, Bruce Nelson², Anthony Freeman ¹, Bruce Chapman¹

1. Jet Propulsion Laboratory California Institute of Technology 4800 Oak Grove Drive Pasadena, California 91109, USA

2. Departamento de Botanica Inst. National de Pesquisas da Amazonia Caixa Postal 478 Manaus AM 69011, Brazil

Abstract

The Amazon basin has the largest continuous region of tropical forest in the world. There has been an increasing interest in the fate of this region as its diversity of species and forest types have been threatened by widespread land-use change and deforestation in the past two decades. There is a general agreement that as the mature tropical forest converts to other land covers, a major part of the terrestrial carbon will be released to the atmosphere through burning, removal of biomass, and decomposition of soil and organic The ambiguity in estimating the terrestrial carbon flux is primarily due to uncertainties in the spatial extent of the various types of primary forest, forest regeneration, and the rate of deforestation. Therefore, mapping and monitoring the Amazon basin have become a priority for several scientific disciplines. In this study, we explore the role of radar remote sensing in mapping forest types in the Amazon basin and estimating and monitoring deforestation. We will make use of a combination of radar sensors such as JPL polarimetric AIRSAR system operating at P-, L-, and C-bands, SIR-C/X-SAR L- and C-band polarimetric system, and JERS- 1 L-band HH polarization The data will be used to understand the capability of SAR imagery in separating and classifying major forest types in the Amazon basin such as lowland moist forest, upper and lower rnontane moist forest, inundated forest (permanent swamp forest, varzea and igapo forests), gallery forest, dry woodlands and savanna, and transitional forests (dominated by bamboo, lianas or palms). Furthermore, the sensitivity of SAR frequency and polarization to forest biomass and structure arc examined in order tocharacterize several clearing practices and forest regeneration within the region. The results indicate that: 1) L-band and P-band HV polarizations show strong sensitivity to secondary biomass regrowth in the early stages of regeneration, 2) HH polarization at these frequencies are suitable for mapping and monitoring the seasonal changes in flooded forests and mangroves, 3) application of C-band channels are limited to nonforested areas. C-band HH polarization can be used to map mangroves in coastal regions. Due to signal saturation, however, it cannot separate the inundated forests throughout the basin accurately. The best channel for large scale mapping and monitoring forest inundation is L-HH, and 4) a combination of SAR backscatter and texture information can be used in classifying dry woodlands and savanna, gallery forests. The examples are given from three experiments over the Amazon basin in Peru and Brazil and the results arc verified by using maps and field data collected during the experiments.

This work has been performed at the Jet I'repulsion Laboratory, California Institute of Technology under a grant from the National Aeronautics and Space Administration.

JET PROPULSION LABORATORY

NOTIFICATION OF CLEARANCE

01/23/97

TO: F. Zee

FROM: L,ogj.sties and Technical information Division

SUBJECT: Notification of Clearance - CL 97-0077

The following title has been cleared by the Document Review Services, Section 644, for public release, presentation, and/or printing in the open literature:

Artificial Neural Network Derived Plant Growth Models

This clearance is issued for an abstract and is valid only for release in the U.S.

Clearance issued by

Mary Sue O'Brien

Document Review Services

Section 644

AUTHORIZATION FOR THE EXTERNAL RELEASE OF INFORMATION

20096 N [CL No. ['] - [']', | (for DRS use only)

Submit with	two copies of the	abstract or full par	<u> </u>	Due Date
nior JPL Author	Section 346	Mail Stop 302-231	Ext. 4-1232	1/14/97
Frank Zee	346		- <u> </u>	
OMPLETE TITLE				
Artificial Neural Network Derived Pla	ant Growth M	Model s	<u></u>	
Foreign X Domestic Accou	unt Code <u>244-20</u>	0014-0-3460 _		Premeeting publication
X ABSTRACT (including extended abstract)				Publication on meeting day
FULL PAPER (including viewgraphs, poster, videocassette)				Postmeeting publication
Journal Nam e	<u>Symposiu</u>	m ⁻ ··		Poster session
Meeting-Subject 12th Man In Space——				Oral presentation
Sponsoring Society Space Research Association				ate June 9-13, 1997
Meeting Location <u>Washington</u>	D.C. 20008	-		
BOOK OR BOOK CHAPTER	FT7		DRS:1	20/97
Assigned Laboratory Task OR	Private V			
X) PUBLICATIONBROCHURE	NEWSLETTER			net outside of NASA
		One		
] Was previously cleared: Clearance No(s)	•			
CLDate CL Date		_ Author(s)		
CL—— — Date	; -	Author(s)		
Nature of this work (p the NASA Advantage of the NASA Advantage o	usly reported in Normation for earlier	ew TechnologyRe	port (NTR) _{No.} .	Delayed or Conditional
FOR SECTION 644 USE ONLY			quantier*	_ _
Editor: —	<u>Ext.</u>	Documen	t No	
Customer Code (RTOP No.)		-	Grou	p Condition
AUTHORIZATION (please use blue ink)	_			
The signatory in this column attests to the technical acc the subject document.	curacy of			
Senior JPL Author Tim Kriebach	1/14/97 Date 1/14/97	Office of F	atents and Nev	w Technology Date
"Manager/Supervisor or equivalent	Date	p:171		1/2 1/22/9 ;
Timothy N. Krabach, Deputy Sect Print Name and Title of Manager/Supervisor NOTE: All full papers require Section or Project Manage All abstracts require Group Supervisor approval	or er approval	Documen	ty Pewiewer	JPL 1330-S R

(over)



AUTHORIZATION FOR THE EXTERNAL RELEASE OF INFORMATION Submit with two copies of the abstract or full paper

Senior JPL Author	Section	Mail Stop	Ext. Due Date			
Frank Zee	346	302-231	4-1232 1/14/97			
COMPLETE TITLE						
Artificial Neural Network Derived Pla	ant Growth Mo	dels	_			
Foreign X Domestic Account Co	ode 244-20014-	0-3460	X Premeeting publication			
ABSTRACT (including extended abstract)			Publication on meeting day			
FULL PAPER (including viewgraphs, poster, vid	Postmeeting publication					
Journal Name	— Poster session					
Title Main Tit by Cooping Control of the Proceedings						
Sponsoring Society Universities Space Research Association Meeting Location Washington, D.C. 20008 Date June 9-13, 1997						
BOOK OR BOOK CHAPTER						
Assigned Laboratory Task OR private Venture						
PUBLICATION BROCHURE NEWS			the Internet outside of NASA			
E FOBLICATION [] BROCHOKE [] NEWS)LL I I L IX					
☐ Was previously cleared: Clearance No(s).:		OKL	RECEIVED			
• • • • • • • • • • • • • • • • • • • •			141 201007			
CL Date		Author(s)				
CL Date		Author(s)	16 TH OFFICE			
REPORTABLE INFORMATION						
THIS WORK: X Not previously reported						
	describe) ADD1s	v i ng artificial	neural networks for			
			rogram			
			(NTR) No			
			(1111) NO			
FOR OFFICE OF PATENTS USE ONLY: Release [-] Ex Post Facto Release Delayed or Conditional						
COMMENTS:						
FOR SECTION 644 USE ONLY						
TOR SECTION 044 USE ONE						
Editor:	Evt	Docum	nent No			
Customer Code (RTOP No.)						
Customer Code (NTOF No.)			Cloup Condition			
AUTHORIZATION (please use blue ink)						
The signatory inthis column attests to the technical accuracy of		0	1.			
the subject document.		. (1)	<u> </u>			
Frank 200 1/14/97 GHB / 1/2/2)						
Senior JPLAuthor, Date Office of Patents and New Technology Date						
Tim Krubach 11	16/97	U				
Manager/Supervisor or equivalent	Date	2				
Timothy .N. Krabach, Deputy Section Ma	inager di	$\mathcal{A}_{\mathcal{K}}$				
Print Name and Title of Manager/Supervisor	īriā Rēr	Document Reviev	ver Date			
NOTE: All /ul/ papers require Section or Project Manager approval. All abstracts require Group Supervisor approval only.						

ARTIFICIAL NEURAL NETWORK DERIVED PLANT GROWTH MODELS *

Frank Zee
Center for Space Microelectronics Technology
Jet Propulsion Laboratory
California Institute of Technology
4800 Oak Grove Dr.
Pasadena, CA 91109-8099

The goal of the Advanced Life Support Systems (ALSS) is to provide self-sufficiency in life support for productive research and exploration in space, for benefits on Earth and to provide a basis for planetary explorations. Part of this objective is to be able to grow crop plants in one or more controlled environments for the purpose of providing life essentials to a human crew, **such as oxygen**, potable water, and food. To do this reliably and **efficiently**, it is necessary to achieve control of the rates of various plant physiology processes. including: net exchange of exhaled carbon dioxide for oxygen (net photosynthesis), purification of water (transpiration), and food production (biomass production rate and harvest index).

To develop an efficient control system that will be able to manage, control, and optimize plant-based life support functions, system identification and modeling of plant growth behavior must first be done. We have developed a plant growth (physiology) model using artificial neural networks. Neural networks are very suitable for both steady-state and dynamic modeling and identification tasks, since they can be trained to approximate arbitrary nonlinear input-output mappings from a collection of input and output examples. In addition, they can be expanded to incorporate a large number of inputs and outputs as required, which makes it simple to model multivariable systems. Thus, unknown nonlinear functions in dynamical models and controllers can easily be parametrized by means of multilayer neural network architectures.

Artificial neural networks are composed of simple albeit numerous non-linear processing elements (modeled after biological neurons) interconnected through a complex network of variable strength connections (modeled after biological synapses). The topology of interconnections and the synaptic strengths essentially dictate the functionality of a given network. A typical network is capable of receiving a large number of analog/digital inputs (e.g., sensor signals) in parallel, and after a complex nonlinear transformation operation, provides the outputs (e.g., predicted growth, biomass). The unique strength of such neural network architectures emerges from their ability to build up their own rules through learning from examples the underlying input/output transformations in ill-defined problems.

In this paper, we will describe our approach to developing these models, the neural network architecture, and the results. With the use of neural networks, these complex, nonlinear, dynamic, multimodal, multivariable plant growth models will be able to better interpolate between all the various environmental conditions and parameters and be able to simulate both short-term (day-to-day) and long-term (plant life cycle) growth of various plants.

^{*}Sponsored by NASA, Office of Life and Microgravity Sciences and Applications